

UTILIZATION MANAGEMENT MEDICAL POLICY

POLICY: Hyperlipidemia – PCSK9 Inhibitors – Leqvio Utilization Management Medical Policy

- Leqvio® (inclisiran subcutaneous injection – Novartis)

REVIEW DATE: 03/04/2026; selected revision 05/06/2026

OVERVIEW

Leqvio, a small interfering ribonucleic acid (RNA) directed to proprotein convertase subtilisin kexin type 9 (PCSK9) messenger RNA, is indicated as an adjunct to diet and exercise to reduce low-density lipoprotein cholesterol (LDL-C) in:¹

- **Hypercholesterolemia** in adults.
- **Heterozygous familial hypercholesterolemia (HeFH)** in adults and pediatric patients ≥ 12 years of age.
- **Homozygous familial hypercholesterolemia (HoFH)** in pediatric patients ≥ 12 years of age

Lerochol™ (lerodalcibep-liga subcutaneous injection), Repatha® (evolocumab subcutaneous injection) and Praluent® (alirocumab subcutaneous injection) are PCSK9 inhibitor products.²⁻⁴

Of note, studies of Leqvio in adults with HoFH did not show reduction in LDL-C levels.⁵

Dosing Information

Leqvio is given as a subcutaneous injection and should be administered by a healthcare professional.¹ The dose is 284 mg given as a single subcutaneous injection initially, again at 3 months, and then once every 6 months.

Guidelines

Several major guidelines address the management of dyslipidemia, including the 2026 American College of Cardiology (ACC)/American Heart Association (AHA) Guideline on the Management of Dyslipidemia, the 2023 ACC/AHA Guideline for the Management of Chronic Coronary Disease (CCD), the 2022 ACC Expert Consensus Decision Pathway (ECPD) on the Role of Non-Statin Therapies, the 2025 ACC/AHA Guideline for the Management of Patients With Acute Coronary Syndrome, the 2026 American Diabetes Association Standards of Care, and the 2025 American Association of Clinical Endocrinology Clinical Practice Guideline for Dyslipidemia.⁶⁻¹¹ Across these guidelines, maximally tolerated statin therapy is consistently identified as the foundation of lipid management, with ezetimibe and/or PCSK9 inhibitors added when additional LDL-C lowering is required. Among PCSK9-targeting therapies, Praluent and Repatha are generally preferred over Leqvio because they have demonstrated cardiovascular (CV) outcomes benefit in large randomized trials, whereas outcomes data for Leqvio are still pending. The guidelines converge in recommending PCSK9 inhibitors for patients with clinical atherosclerotic cardiovascular disease (ASCVD) or severe hypercholesterolemia (LDL-C ≥ 190 mg/dL) who remain above LDL-C goals despite optimized background therapy; however, they differ in how treatment targets are expressed. The 2026 ACC/AHA dyslipidemia guideline and the 2026 ADA Standards of Care define explicit LDL-C goals, including a target of < 55 mg/dL for patients with very high-risk ASCVD, whereas other guidance, such as the 2022 ACC Expert Consensus Decision Pathway and the 2023 Chronic Coronary Disease guideline, relies primarily on LDL-C thresholds to trigger treatment intensification rather than strict numeric goals.

The AHA Scientific Statement on Familial Hypercholesterolemia (2015) and other sources provide guidance on the diagnosis of familial hypercholesterolemia, including HeFH.^{12,13} Diagnostic approaches include the Dutch Lipid Clinic Network scoring system and the Simon Broome criteria.

POLICY STATEMENT

Prior Authorization is recommended for medical benefit coverage of Leqvio. Approval is recommended for those who meet the **Criteria** and **Dosing** for the listed indications. Extended approvals are allowed if the patient continues to meet the Criteria and Dosing. Requests for doses outside of the established dosing documented in this policy will be considered on a case-by-case basis by a clinician (i.e., Medical Director or Pharmacist). All approvals are provided for the duration noted below. A patient who has previously met Initial Therapy criteria for Leqvio for the requested indication under the Coverage Review Department and is currently receiving Leqvio is only required to meet continuation of therapy criteria (i.e., currently receiving therapy). If past criteria have not been met under the Coverage Review Department and the patient is currently receiving Leqvio, or is restarting Leqvio, Initial Therapy criteria must be met.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Leqvio is recommended in those who meet one of the following criteria:

FDA-Approved Indications

1. **Heterozygous Familial Hypercholesterolemia (HeFH).*** Approve for 1 year if the patient meets ONE of the following (A or B):
 - A) **Initial Therapy.** Approve if the patient meets ALL of the following (i, ii, and iii):
 - i. Patient is ≥ 12 years of age; AND
 - ii. Patient meets ONE of the following (a, b, c, or d):
 - a) Patient has an untreated low-density lipoprotein cholesterol (LDL-C) level ≥ 190 mg/dL (prior to treatment with antihyperlipidemic agents); OR
 - b) If the patient is between 12 and 17 years of age, meets BOTH of the following [(1) and (2)]:
 - (1) Patient has an untreated low-density lipoprotein cholesterol (LDL-C) ≥ 160 mg/dL (prior to treatment with antihyperlipidemia agents); AND
 - (2) According to the prescriber, patient has a family history of early atherosclerotic cardiovascular disease (ASCVD) or elevated low-density lipoprotein cholesterol (LDL-C) or total cholesterol (TC) in a parent; OR
 - c) The diagnosis has been confirmed by genetic testing; OR
Note: Examples include pathogenic variants at the low-density lipoprotein receptor (LDLR), apolipoprotein B (APOB), proprotein convertase subtilisin kexin type 9 (PCSK9), or low-density lipoprotein receptor adaptor protein 1 (LDLRAP1) gene.
 - d) Patient has been diagnosed with heterozygous familial hypercholesterolemia meeting ONE of the following diagnostic criteria thresholds [(1) or (2)]:
 - (1) Prescriber confirms that the Dutch Lipid Network criteria score was > 5 ; OR
 - (2) Prescriber confirms that Simon Broome criteria met the threshold for “definite” or “possible (or probable)” familial hypercholesterolemia; AND
 - iii. Patient meets ONE of the following (a or b):
 - a) Patient meets BOTH of the following [(1) and (2)]:

- (1) Patient has tried one high-intensity statin therapy (i.e., atorvastatin $\geq$ 40 mg daily; rosuvastatin $\geq$ 20 mg daily [as a single entity or as a combination product]) for $\geq$ 8 continuous weeks; AND
- (2) LDL-C level after this treatment remains $\geq$ 70 mg/dL; OR
- b)** Patient has been determined to be statin-intolerant by meeting ONE of the following [(1) or (2)]:
 - (1) Patient experienced statin-related rhabdomyolysis; OR
Note: Rhabdomyolysis is a statin-induced muscle breakdown that is associated with markedly elevated creatine kinase levels (at least 10 times the upper limit of normal), along with evidence of end organ damage which can include signs of acute renal injury (noted by substantial increases in serum creatinine [Scr] levels [$\geq$ 0.5 mg/dL increase in Scr or doubling of the Scr] and/or myoglobinuria [myoglobin present in urine]).
 - (2) Patient meets ALL of the following [(a), (b), and (c)]:
 - (a) Patient experienced skeletal-related muscle symptoms; AND
Note: Examples of skeletal-related muscle symptoms include myopathy (muscle weakness) and myalgia (muscle aches, soreness, stiffness, or tenderness).
 - (b) The skeletal-muscle related symptoms occurred while receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or combination products); AND
 - (c) When receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or as combination products) the skeletal-related muscle symptoms resolved upon discontinuation of each respective statin therapy (atorvastatin and rosuvastatin); OR
Note: Examples of skeletal-related muscle symptoms include myopathy and myalgia.
- B)** Patient is Currently Receiving Leqvio. Approve if according to the prescriber, the patient has experienced a response to therapy.
Note: Examples of a response to therapy include decreasing LDL-C, total cholesterol, non-high-density lipoprotein (non-HDL-C), or apolipoprotein B levels. Also, if the patient is currently receiving the requested therapy but has not previously received approval of Leqvio for this specific indication through the Coverage Review Department, review under criteria for Initial Therapy. If the patient is restarting therapy with Leqvio, Initial Therapy criteria must be met.

Dosing. Approve ONE of the following dosage regimens (A or B):

- A)** Initial dose is 284 mg given as a single subcutaneous injection, again at 3 months, and then once every 6 months; OR
- B)** Maintenance dose is 284 mg given as a subcutaneous injection once every 6 months.

2. Homozygous Familial Hypercholesterolemia (HoFH).* Approve for 1 year if the patient meets ONE of the following (A or B):

- A)** Initial Therapy. Approve if the patient meets ALL of the following (i, ii, and iii):
 - i.** Patient is $\geq$ 12 years of age and $<$ 18 years of age; AND
 - ii.** Patient meets ONE of the following (a, b, or c):
 - a)** The diagnosis has been confirmed by genetic testing; OR
Note: Examples include pathogenic variants at the low-density lipoprotein receptor (LDLR), apolipoprotein B (APOB), proprotein convertase subtilisin kexin type 9 (PCSK9) or low-density lipoprotein receptor adaptor protein (LDLRAP1) gene.
 - b)** Patient has an untreated low-density lipoprotein (LDL-C) level $>$ 400 mg/dL AND meets ONE of the following [(1) or (2)]:
Note: Untreated refers to prior therapy with any antihyperlipidemic agent.

- (1) Patient had clinical manifestations of homozygous familial hypercholesterolemia before 10 years of age; OR
Note: Clinical manifestations of homozygous familial hypercholesterolemia are cutaneous xanthomas, tendon xanthomas, arcus cornea, tuberous xanthomas, or xanthelasma.
- (2) At least one parent of the patient had untreated LDL-C levels or total cholesterol levels consistent with familial hypercholesterolemia; OR
Note: An example of familial hypercholesterolemia is an untreated LDL-C level ≥ 190 mg/dL and/or an untreated total cholesterol level > 250 mg/dL.
- c) Patient has a treated LDL-C level ≥ 300 mg/dL AND meets ONE of the following [(1) or (2)]:
Note: Treated refers to after therapy with at least one antihyperlipidemic agent. Some examples of antihyperlipidemic agents include statins (e.g., atorvastatin, rosuvastatin, lovastatin, simvastatin, pravastatin), ezetimibe, a PCSK9 inhibitor (e.g., Repatha [evolocumab subcutaneous injection]), Evkeeza (evinacumab-dgnb intravenous infusion), and Juxtapid (lomitapide capsules).
 - (1) Patient had clinical manifestations of homozygous familial hypercholesterolemia before 10 years of age; OR
Note: Examples of clinical manifestations of homozygous familial hypercholesterolemia are cutaneous xanthomas, tendon xanthomas, arcus cornea, tuberous xanthomas or xanthelasma.
 - (2) At least one parent of the patient had untreated LDL-C levels or total cholesterol levels consistent with familial hypercholesterolemia; AND
Note: An example of familial hypercholesterolemia is an untreated LDL-C ≥ 190 mg/dL and/or an untreated total cholesterol > 250 mg/dL.
- iii. Patient meets ONE of the following (a or b):
 - a) Patient meets BOTH of the following [(1) and (2)]:
 - (1) Patient has tried one high-intensity statin therapy (i.e., atorvastatin ≥ 40 mg daily; rosuvastatin ≥ 20 mg daily [as a single-entity or as a combination product]) for ≥ 8 continuous weeks; AND
 - (2) LDL-C level after this treatment remains ≥ 70 mg/dL; OR
 - b) Patient has been determined to be statin-intolerant by meeting ONE of the following [(1) or (2)]:
 - (1) Patient experienced statin-related rhabdomyolysis; OR
Note: Rhabdomyolysis is a statin-induced muscle breakdown that is associated with markedly elevated creatine kinase levels (at least 10 times the upper limit of normal), along with evidence of end organ damage which can include signs of acute renal injury (noted by substantial increases in serum creatinine [Scr] levels [a ≥ 0.5 mg/dL increase in Scr or doubling of the Scr] and/or myoglobinuria [myoglobin present in urine]); OR
 - (2) Patient meets ALL of the following [(a), (b), and (c)]:
 - (a) Patient experienced skeletal-related muscle symptoms; AND
Note: Examples of skeletal-related muscle symptoms include myopathy (muscle weakness) and myalgia (muscle aches, soreness, stiffness, or tenderness).
 - (b) The skeletal-muscle related symptoms occurred while receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or combination products); AND
 - (c) When receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or as combination products) the skeletal-related muscle symptoms resolved upon discontinuation of each respective statin therapy (atorvastatin and rosuvastatin); OR
Note: Examples of skeletal-related muscle symptoms include myopathy and myalgia.

- B) Patient is Currently Receiving Leqvio.** Approve if according to the prescriber, the patient has experienced a response to therapy.

Note: Examples of a response to therapy include decreasing low-density lipoprotein cholesterol (LDL-C), total cholesterol, non-high-density lipoprotein (non-HDL-C), or apolipoprotein B levels. Also, if the patient is currently receiving the requested therapy but has not previously received approval of Leqvio for this specific indication through the Coverage Review Department, review under criteria for Initial Therapy. If the patient is restarting therapy with Leqvio, Initial Therapy criteria must be met.

Dosing. Approve ONE of the following dosage regimens (A or B):

- A)** Initial dose is 284 mg given as a single subcutaneous injection, again at 3 months, and then once every 6 months; OR
- B)** Maintenance dose is 284 mg given as a subcutaneous injection once every 6 months.

3. Hypercholesterolemia.* Approve for 1 year if the patient meets ONE of the following (A or B):

Note: This is not associated with established cardiovascular disease or heterozygous familial hypercholesterolemia (HeFH) and may be referred to as combined hyperlipidemia, hypercholesterolemia (pure, primary), dyslipidemia, or increased/elevated low-density lipoprotein cholesterol (LDL-C) levels.

- A) Initial Therapy.** Approve if the patient meets ALL of the following (i, ii, and iii):

i. Patient is ≥ 18 years of age; AND

ii. Patient meets ONE of the following (a or b):

a) According to the prescriber, the patient is at elevated risk for atherosclerotic cardiovascular disease (ASCVD), an ASCVD event, or a major adverse cardiovascular event (MACE); OR

b) Patient has diabetes; AND

iii. Patient meets ONE of the following (a or b):

a) Patient meets ALL of the following [(1) and (2)]:

(1) Patient has tried one high-intensity statin therapy (i.e., atorvastatin ≥ 40 mg daily; rosuvastatin ≥ 20 mg daily [as a single-entity or as a combination product]); AND

(2) LDL-C level after this treatment regimen remains ≥ 70 mg/dL; OR

b) Patient has been determined to be statin-intolerant by meeting ONE of the following [(1) or (2)]:

(1) Patient experienced statin-related rhabdomyolysis; OR

Note: Rhabdomyolysis is a statin-induced muscle breakdown that is associated with markedly elevated creatine kinase levels (at least 10 times the upper limit of normal), along with evidence of end organ damage, which can include signs of acute renal injury (noted by substantial increases in serum creatinine [Scr] levels [$a \geq 0.5$ mg/dL increase in Scr or doubling of the Scr] and/or myoglobinuria [myoglobin present in urine]).

(2) Patient meets ALL of the following [(a), (b), and (c)]:

(a) Patient experienced skeletal-related muscle symptoms; AND

Note: Examples of skeletal-related muscle symptoms include myopathy (muscle weakness) and myalgia (muscle aches, soreness, stiffness, or tenderness).

(b) The skeletal-muscle related symptoms occurred while receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or combination products); AND

(c) When receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or as combination products), the skeletal-related muscle symptoms resolved upon discontinuation of each respective statin therapy (atorvastatin and rosuvastatin); OR

Note: Examples of skeletal-related muscle symptoms include myopathy and myalgia.

- B) Patient is Currently Receiving Leqvio.** Approve if according to the prescriber, the patient has experienced a response to therapy.

Note: Examples of a response to therapy include decreasing LDL-C, total cholesterol, non-high-density lipoprotein (non-HDL-C), or apolipoprotein B levels. Also, if the patient is currently receiving the requested therapy but has not previously received approval of Leqvio for this specific indication through the Coverage Review Department, review under criteria for Initial Therapy. If the patient is restarting therapy with Leqvio, Initial Therapy criteria must be met.

Dosing. Approve ONE of the following dosage regimens (A or B):

- A)** Initial dose is 284 mg given as a single subcutaneous injection, again at 3 months, and then once every 6 months; OR
- B)** Maintenance dose is 284 mg given as a subcutaneous injection once every 6 months.

Other Uses with Supportive Evidence

- 3. Established Cardiovascular Disease.*** Approve for 1 year if the patient meets ONE of the following (A or B):

- A) Initial Therapy.** Approve if the patient meets ALL of the following (i, ii, and iii):

- i.** Patient is ≥ 18 years of age; AND
- ii.** Patient has had ONE of the following conditions or diagnoses (a, b, c, d, e, or f):
 - a)** A previous myocardial infarction or a history of an acute coronary syndrome; OR
 - b)** Angina (stable or unstable); OR
 - c)** A past history of stroke or transient ischemic attack; OR
 - d)** Coronary artery disease; OR
 - e)** Peripheral arterial disease; OR
 - f)** Patient has undergone a coronary or other arterial revascularization procedure in the past; AND

Note: Examples include coronary artery bypass graft surgery, percutaneous coronary intervention, angioplasty, and coronary stent procedures.

- iii.** Patient meets ONE of the following (a or b):

- a)** Patient meets BOTH of the following [(1) and (2)]:
 - (1)** Patient has tried one high-intensity statin therapy (i.e., atorvastatin ≥ 40 mg daily; rosuvastatin ≥ 20 mg daily [as a single entity or as a combination product]) for ≥ 8 continuous weeks; AND
 - (2)** Low-density lipoprotein cholesterol (LDL-C) level after this treatment remains ≥ 55 mg/dL; OR
- b)** Patient has been determined to be statin-intolerant by meeting ONE of the following [(1) or (2)]:
 - (1)** Patient experienced statin-related rhabdomyolysis; OR

Note: Rhabdomyolysis is a statin-induced muscle breakdown that is associated with markedly elevated creatine kinase levels (at least 10 times the upper limit of normal), along with evidence of end organ damage which can include signs of acute renal injury (noted by substantial increases in serum creatinine [Scr] levels [a ≥ 0.5 mg/dL increase in Scr or doubling of the Scr] and/or myoglobinuria [myoglobin present in urine]).
 - (2)** Patient meets ALL of the following [(a), (b), and (c)]:
 - (a)** Patient experienced skeletal-related muscle symptoms; AND

Note: Examples of skeletal-related muscle symptoms include myopathy (muscle weakness) and myalgia (muscle aches, soreness, stiffness, or tenderness).

- (b) The skeletal-muscle related symptoms occurred while receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or combination products); AND
- (c) When receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or as combination products) the skeletal-related muscle symptoms resolved upon discontinuation of each respective statin therapy (atorvastatin and rosuvastatin); OR

Note: Examples of skeletal-related muscle symptoms include myopathy and myalgia.

- B) Patient is Currently Receiving Leqvio.** Approve if according to the prescriber, the patient has experienced a response to therapy.

Note: Examples of a response to therapy include decreasing LDL-C, total cholesterol, non-high-density lipoprotein (non-HDL-C), or apolipoprotein B levels. Also, if the patient is currently receiving the requested therapy but has not previously received approval of Leqvio for this specific indication through the Coverage Review Department, review under criteria for Initial Therapy. If the patient is restarting therapy with Leqvio, Initial Therapy criteria must be met.

Dosing. Approve ONE of the following dosage regimens (A or B):

- C)** Initial dose is 284 mg given as a single subcutaneous injection, again at 3 months, and then once every 6 months; OR
- D)** Maintenance dose is 284 mg given as a subcutaneous injection once every 6 months.

Note:

* A patient may have a diagnosis that pertains to more than one indication, therefore, consider review under different approval conditions, if applicable (e.g., a patient with heterozygous familial hypercholesterolemia may have established cardiovascular disease, a patient with primary hyperlipidemia may have heterozygous familial hypercholesterolemia).

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Leqvio is not recommended in the following situations:

- 1. Concurrent use of Leqvio with Lerochol (lerodalcibep-liga subcutaneous injection), Repatha (evolocumab subcutaneous injection) or Praluent (alirocumab subcutaneous injection).** Lerochol, Repatha, and Praluent are PCSK9 inhibitors and should not be used with Leqvio due to a similar mechanism of action.¹ Patients receiving PCSK9 inhibitors were excluded from the pivotal trials with Leqvio.
- 2. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria.** Criteria will be updated as new published data are available.

REFERENCES

1. Leqvio® subcutaneous injection [prescribing information]. East Hanover, NJ: Novartis; February 2026.
2. Lerochol™ subcutaneous injection [prescribing information]. Cincinnati, OH: LIB; December 2025.
3. Repatha® subcutaneous injection [prescribing information]. Thousand Oaks, CA: Amgen; August 2025.
4. Praluent® subcutaneous injection [prescribing information]. Tarrytown, NY: Regeneron; October 2025.
5. Raal F, Durst R, Bi R, et al. Efficacy, safety, and tolerability of inclisiran in patients with homozygous familial hypercholesterolemia: results from the ORION-5. *Circulation*. 2024;149:354-362.
6. Lloyd-Jones DM, Morris PB, Ballantyne CM, et al. 2022 ACC Expert Consensus Decision Pathway on the Role of Non-Statins Therapies for LDL-Cholesterol Lowering in the Management of Atherosclerotic Cardiovascular Disease Risk. *J Am Coll Cardiol*. 2022;80(14):1366-1418.

7. Blumenthal RS, Morris PB, Gaudino M, et al. 2026 ACC/AHA/AACVPR/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA guideline on the management of dyslipidemia: a report of the American College of Cardiology/American Heart Association joint committee on clinical practice guidelines. *JACC*. 2026 March 13. [Online ahead of print].
8. American Diabetes Association Professional Practice Committee. Cardiovascular Disease and Risk Management: Standards of Care in Diabetes-2026. *Diabetes Care*. 2026;49(Suppl 1):S216-S245.
9. Virani SS, Newby LK, Arnold SV, et al. 2023 AHA/ACC/ACCP/ASPC/NLA/PCNA guideline for the management of patients with chronic coronary disease: a report of the American Heart Association/American College of Cardiology Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2023;82(9):833-955.
10. Rao SV, O'Donoghue ML, Ruel M, et al. 2025 ACC/AHA/ACEP/NAEMSP/SCAI guideline for the management of patients with acute coronary syndromes. *J Am Coll Cardiol*. 2025 Feb 27. [Online ahead of print].
11. Patel SB, Wyne KL, Afreen S, et al. American Association of Clinical Endocrinology clinical practice guideline on pharmacologic management of adults with dyslipidemia. *Endocrine Pract*. 2025;31:236-262.
12. Gidding SS, Champagne MA, de Ferranti SD, et al. The agenda for familial hypercholesterolemia. A scientific statement from the American Heart Association. *Circulation*. 2015;132(22):2167-2192.
13. Haase A, Goldberg AC. Identification of people with heterozygous familial hypercholesterolemia. *Curr Opin Lipidol*. 2012;23:282-289.

HISTORY

Type of Revision	Summary of Changes	Review Date
Annual Revision	It was removed from the Policy Statement that the agent is prescribing by or in consultation with a physician who specializes in the condition being treated. In addition, the following changes were made: Established Cardiovascular Disease: The name of the indication was changed to as stated (previously “Atherosclerotic Cardiovascular Disease”). For <u>Initial Therapy</u>, the requirement that the medication is prescribed by, or in consultation with a cardiologist; an endocrinologist; or a physician who focuses in the treatment of cardiovascular risk management and/or lipid disorders was removed. The requirement that the low-density lipoprotein cholesterol level after treatment with one high-intensity statin therapy and ezetimibe be ≥ 70 mg/dL was changed to ≥ 55 mg/dL. For a <u>Patient Currently Receiving the Medication</u>, the requirement that the “prescribing physician” notes that the patient has experienced a response to therapy was changed to “prescriber”. Heterozygous Familial Hypercholesterolemia: For <u>Initial Therapy</u>, the requirement that the medication is prescribed by, or in consultation with a cardiologist; an endocrinologist; or a physician who focuses in the treatment of cardiovascular risk management and/or lipid disorders was removed. The requirement that the patient has had genetic confirmation of heterozygous familial hypercholesterolemia by mutations in the low-density lipoprotein receptor, apolipoprotein B, proprotein convertase subtilisin kexin type 9, or low-density lipoprotein receptor adaptor protein 1 gene was changed to state that the patient has had phenotypic confirmation of heterozygous familial hypercholesterolemia with the above examples moved to a Note. Regarding the diagnosis of heterozygous familial hypercholesterolemia by meeting the Dutch Lipid Network criteria score or the Simon Broome criteria, the requirement that this be confirmed by the “prescribing physician” was changed to “prescriber”. For a <u>Patient Currently Receiving the Medication</u>, the requirement that the “prescribing physician” notes that the patient has experienced a response to therapy was changed to “prescriber”. Primary Hyperlipidemia: For <u>Initial Therapy</u>, the requirement that the medication is prescribed by, or in consultation with a cardiologist; an endocrinologist; or a physician who focuses in the treatment of cardiovascular risk management and/or lipid disorders was removed. A patient with diabetes now qualifies for this indication (if requirements are met); previously, high risk was only defined by a patient who had a “coronary artery calcium or calcification score ≥ 300 Agatston units”. The requirement that the low-density lipoprotein cholesterol level after treatment with one high-intensity statin therapy, along with ezetimibe, be ≥ 100 mg/dL was changed to ≥ 70 mg/dL. For a <u>Patient Currently Receiving the Medication</u>, the requirement that the “prescribing physician” notes that the patient has experienced a response to therapy was changed to “prescriber”.	05/08/2024
Annual Revision	Heterozygous Familial Hypercholesterolemia (HeFH): For <u>Initial Therapy</u>, the phrase “phenotypic confirmation of heterozygous familial hypercholesterolemia” was replaced with “The diagnosis has been confirmed by genetic testing”. Also, “apo B” was changed to “APOB”.	05/28/2025

HISTORY (CONTINUED)

Type of Revision	Summary of Changes	Review Date
Selected Revision	Heterozygous Familial Hypercholesterolemia: For initial therapy the requirement that the patient has tried one high-intensity statin along with ezetimibe (as a single-entity or as a combination product) for ≥ 8 continuous weeks was removed. The requirement remains that the patient has tried one high-intensity statin therapy (i.e., atorvastatin ≥ 40 mg daily; rosuvastatin ≥ 20 mg daily [as a single entity or as a combination product]) and the qualifier of “for ≥ 8 continuous weeks” was added for clarification. Established Cardiovascular Disease: For initial therapy the requirement that the patient has tried one high-intensity statin along with ezetimibe (as a single-entity or as a combination product) for ≥ 8 continuous weeks was removed. The requirement remains that the patient has tried one high-intensity statin therapy (i.e., atorvastatin ≥ 40 mg daily; rosuvastatin ≥ 20 mg daily [as a single entity or as a combination product]) and the qualifier of “for ≥ 8 continuous weeks” was added for clarification.	08/06/2025
Update	The policy name was changed to as listed. Previously, it was Proprotein Convertase Subtilisin Kexin Type 9 Related Products – Leqvio UM Medical Policy.	N/A
Early Annual Revision	Heterozygous Familial Hypercholesterolemia: The age requirement was changed to approve for a patient ≥ 12 years of age; previously was ≥ 18 years of age. For a patient between 12 and 17 years of age, a requirement was added that untreated low-density lipoprotein cholesterol (LDL-C) is ≥ 160 mg/dL prior to treatment with antihyperlipidemic agents, and that, according to the prescriber, the patient has a family history of early atherosclerotic cardiovascular disease or elevated LDL-C or total cholesterol in a parent. Homozygous Familial Hypercholesterolemia: This was added as a new condition of approval. Hypercholesterolemia: The diagnosis of Primary Hyperlipidemia was changed to as listed. A Note was also updated with this reworded indication. Conditions Not Recommended for Approval: Lerochol was added as an agent that cannot be taken concurrently with Leqvio.	03/04/2026
Selected Revision	Hypercholesterolemia: The requirement for a patient to have a coronary artery calcium or calcification score ≥ 300 was replaced with a requirement that, according to the prescriber, the patient is at risk for atherosclerotic cardiovascular disease (ASCVD), an ASCVD event, or a major adverse cardiovascular event (MACE). The requirement for a patient to have tried one high-intensity statin therapy along with ezetimibe (as a single-entity or as a combination product) for ≥ 8 continuous weeks was removed.	05/06/2026

APPENDIX A**Simon Broome Register Diagnostic Criteria.^{12,13}**

Definite Familial Hypercholesterolemia
Raised cholesterol
--Total cholesterol greater than 6.7 mmol/L (260 mg/dL) or LDL-C > 4.0 mmol/L (155 mg/dL) in a patient < 16 years of age; OR
--Total cholesterol > 7.5 mmol/L (290 mg/dL) or LDL-C > 4.9 mmol/L (190 mg/dL) in a patient > 16 years of age;
AND
--Tendon xanthomas in the patient or in a first (parent, sibling, or child) or second-degree relative (grandparent, aunt, or uncle);
OR
DNA-based evidence of LDL-receptor, familial defective APOB, or PCSK9 mutation.
Possible (or Probable) Familial Hypercholesterolemia
Raised cholesterol
--Total cholesterol greater than 6.7 mmol/L (260 mg/dL) or LDL-C > 4.0 mmol/L (155 mg/dL) in a patient < 16 years of age; OR
--Total cholesterol > 7.5 mmol/L (290 mg/dL) or LDL-C > 4.9 mmol/L (190 mg/dL) in a patient > 16 years of age;
AND
Family history of premature myocardial infarction younger than 50 years of age in second-degree relative or younger than 60 years of age in first-degree relative;
OR
Raised cholesterol
--Total cholesterol greater than 6.7 mmol/L (260 mg/dL) or LDL-C > 4.0 mmol/L (155 mg/dL) in a patient < 16 years of age; OR
--Total cholesterol > 7.5 mmol/L (290 mg/dL) or LDL-C > 4.9 mmol/L (190 mg/dL) in a patient > 16 years of age;
AND
Family history of raised cholesterol > 7.5 mmol (290 mg/dL) in adult first-degree or second-degree relative or > 6.7 mmol/L (260 mg/dL) in child or sibling aged < 16 years.

LDL-C – Low-density lipoprotein cholesterol; LDL – Low-density lipoprotein; APOB – Apolipoprotein B; PCSK9 – Proprotein convertase subtilisin kexin type 9.

APPENDIX B.

Dutch Lipid Network Criteria.^{12,13}

Criteria	Score
Family History	
First-degree relative with known premature coronary and/or vascular disease (men < 55 years, women < 60 years)	1
First degree relative with known LDL-C > 95 th percentile for age and sex	1
First-degree relative with tendon xanthomata and/or arcus cornealis, OR	2
Patient is < 18 years of age with LDL-C > 95 th percentile for age and sex	2
Clinical History	
Patient with premature CAD (age as above)	2
Patient with premature cerebral or peripheral vascular disease (age as above)	1
Physical Examination	
Tendon xanthomas	6
Arcus cornealis at age < 45 years	4
LDL-C	
LDL-C ≥ 8.5 mmol/L (330 mg/dL)	8
LDL-C 6.5 to 8.4 mmol/L (250 to 329 mg/dL)	5
LDL-C 5.0 to 6.4 mmol/L (190 to 249 mg/dL)	3
LDL-C 4.0 to 4.9 mmol/L (155 to 189 mg/dL)	1
DNA Analysis	
Functional mutation LDLR, APOB or PCSK9 gene	8
Stratification	
Definite familial hypercholesterolemia	> 8
Probable familial hypercholesterolemia	6 to 8
Possible familial hypercholesterolemia	3 to 5
Unlikely familial hypercholesterolemia	< 3

LDL-C – Low-density lipoprotein cholesterol; CAD – Coronary artery disease; LDLR – Low-density lipoprotein receptor; APOB – Apolipoprotein B; PCSK9 – Proprotein convertase subtilisin kexin type 9.